

Views About Management



A statement of English Nature's views about the management of Prior's Park & Adcombe Woods Site of Special Scientific Interest (SSSI).

This statement represents English Nature's views about the management of the SSSI for nature conservation. This statement sets out, in principle, our views on how the site's special conservation interest can be conserved and enhanced. English Nature has a duty to notify the owners and occupiers of the SSSI of its views about the management of the land.

Not all of the management principles will be equally appropriate to all parts of the SSSI. Also, there may be other management activities, additional to our current views, which can be beneficial to the conservation and enhancement of the features of interest.

The management views set out below do not constitute consent for any operation. English Nature's written consent is still required before carrying out any operation likely to damage the features of special interest (see your SSSI notification papers for a list of these operations). English Nature welcomes consultation with owners, occupiers and users of the SSSI to ensure that the management of this site conserves and enhances the features of interest, and to ensure that all necessary prior consents are obtained.

Management Principles

Broadleaved semi-natural woodland

There are many different ways in which broadleaved woodland can be managed to conserve its value for wildlife. The following gives broad views on a range of regimes that may be appropriate on your site.

A diverse woodland structure, with open space, a dense understory, and a more mature overstory is important. A range of ages and species within and between stands is desirable. Some dead and decaying wood, such as fallen logs, can provide habitats for fungi and invertebrates. However, work may be needed to make safe dangerous trees in areas of high public access. Both temporary and permanent open spaces benefit groups of invertebrates such as butterflies. They may require cutting to keep them open, and should be of sufficient size to ensure that sunny conditions prevail for most of the day.

Felling, thinning or coppicing may be used to create or maintain variations in the structure of the wood, and non-native trees and shrubs can be removed at this time. To avoid disturbance to breeding birds and dormice the work is normally best done between the beginning of October and the end of February. Work should be avoided when the ground is soft, to prevent disturbing the soil and ground flora. Normally

successive felling, thinning or coppicing operations should be spread through the wood to promote diversity, but where there is open space adjacent plots should be worked to encourage the spread of species that are only weakly mobile. Natural regeneration from seed or stump regrowth is preferred to planting because it helps maintain the local patterns of species and the inherent genetic character of the site.

Deer management and protection from rabbits or livestock are often necessary. Whilst light or intermittent grazing may increase woodland diversity, heavy browsing can damage the ground flora and prevent successful regeneration. Invasive species, such as *Rhododendron* or Himalayan balsam, should be controlled.

Parts of a wood should be left unmanaged to benefit species that do best under low disturbance or in response to natural processes. Within these areas some trees will eventually die naturally and dead wood accumulate.

Marshy grassland

Marshy grassland requires active management if it is to retain its conservation interest. Generally, each year's growth of vegetation must be removed. Otherwise the sward becomes dominated by tall, vigorous grasses and rushes which, together with an associated build up of dead plant matter, suppress less vigorous species and lower the botanical richness of the sward. Traditionally, this management is achieved by grazing. Cattle are often the preferred stock, being relatively tolerant of wet conditions and able to control tall grasses and rank vegetation. Cattle also tend to produce a rather uneven, structurally diverse sward. However, ponies, or even hill sheep, can be used if necessary. Grazing usually takes place at times between late spring and early autumn, but the precise timing and intensity will depend on local conditions and requirements, such as the need to avoid trampling ground-nesting birds. Heavy poaching should be avoided but light trampling can be beneficial in breaking down leaf litter and providing areas for seed germination. An element of managed scrub, both within and fringing a field can be of importance to birds and invertebrates, as can a surrounding hedge. Careful maintenance of existing ditches and drains is usually acceptable practice, but abandonment or deepening of ditches can be harmful.

All habitats

The habitats within this site are highly sensitive to inorganic fertilisers and pesticides, applications of which should be avoided both within the site itself and in adjacent surrounding areas. Herbicides may be useful in targeting certain invasive species, but should be used with extreme care. Access to this site, and any recreational activities within, may also need to be controlled.