

Views About Management



A statement of English Nature's views about the management of Lount Meadows Site of Special Scientific Interest (SSSI).

This statement represents English Nature's views about the management of the SSSI for nature conservation. This statement sets out, in principle, our views on how the site's special conservation interest can be conserved and enhanced. English Nature has a duty to notify the owners and occupiers of SSSI of its views about the management of the land.

Not all of the management principles will be equally appropriate to all parts of the SSSI. Also, there may be other management activities, additional to our current views, which can be beneficial to the conservation and enhancement of the features of interest.

The management views set out below do not constitute consent for any operation. English Nature's written consent is still required before carrying out any operation likely to damage the features of special interest (see your SSSI notification papers for a list of these operations). English Nature welcomes consultation with owners, occupiers and users of the SSSI to ensure that the management of this site conserves and enhances the features of interest, and to ensure that all necessary prior consents are obtained.

Management Principles

Neutral hay meadows require active management if they are to retain their conservation interest. In order to maintain a species-rich sward, each year's growth of vegetation must be removed. Otherwise the sward becomes progressively dominated by tall and vigorous grasses which, together with an associated build up of dead plant matter, suppress less vigorous species and reduce the botanical diversity of the site. In neutral hay meadows, the above objective is traditionally achieved by closing the fields to stock in the autumn and cutting the resultant growth as hay, usually in mid-July. The precise timing of the cut depends on local factors, including past management and current weather conditions, but should be after ground-nesting birds have fledged their young and any short-lived, characteristic plants have set seed. The aftermath is then grazed in late summer/autumn. Aftermath grazing is important for maintaining a species-rich sward, both through controlling competitive grasses and through hoof-prints providing suitable sites for seedlings to establish. Heavy poaching must be avoided, however. Any surrounding, well managed hedgerows may considerably add to the habitat in providing shelter for invertebrates. The application of pesticides including herbicides or fertilizers would be damaging but periodic dressings of well-rotted farmyard manure may be acceptable if the sward does not receive regular input of nutrients from flooding. Occasional dressings of lime may be acceptable.

For the damper meadows, regular and careful maintenance of surface drainage including ditches and drains can be necessary to prevent adverse changes in the plant species composition of the sward. Deepening of surface drainage should be avoided.