

Views About Management

A statement of English Nature's views about the management of Ge-Mare Farm Fields Site of Special Scientific Interest (SSSI).

This statement represents English Nature's views about the management of the SSSI for nature conservation. This statement sets out, in principle, our views on how the site's special conservation interest can be conserved and enhanced. English Nature has a duty to notify the owners and occupiers of the SSSI of its views about the management of the land.

Not all of the management principles will be equally appropriate to all parts of the SSSI. Also, there may be other management activities, additional to our current views, which can be beneficial to the conservation and enhancement of the features of interest.

The management views set out below do not constitute consent for any operation. English Nature's written consent is still required before carrying out any operation likely to damage the features of special interest (see your SSSI notification papers for a list of these operations). English Nature welcomes consultation with owners, occupiers and users of the SSSI to ensure that the management of this site conserves and enhances the features of interest, and to ensure that all necessary prior consents are obtained.

Management Principles

Marshy grassland

Wet grassland requires active management if it is to retain its conservation interest. Generally, each year's growth of vegetation must be removed. Otherwise the sward becomes dominated by tall, vigorous grasses and rushes which, together with an associated build up of dead plant matter, suppress less vigorous species and lower the botanical richness of the sward. Traditionally, this management is achieved by grazing. Cattle are often the preferred stock, being relatively tolerant of wet conditions and able to control tall grasses and rank vegetation. Cattle also tend to produce a rather uneven, structurally diverse sward. However, ponies, or even hill sheep, can be used if necessary. Grazing usually takes place at times between late spring and early autumn, but the precise timing and intensity will depend on local conditions and requirements, such as the need to avoid trampling ground-nesting birds. Heavy poaching should be avoided but light trampling can be beneficial in breaking down leaf litter and providing areas for seed germination. An element of managed scrub, both within and fringing a field can be of importance to birds and invertebrates, as can a surrounding hedge. Careful maintenance of existing ditches and drains is usually acceptable practice, but abandonment or deepening of ditches can be harmful. Cultivation, increased drainage or the application of pesticides, including herbicides, or fertilizer is likely to be damaging and should be avoided.