

Views About Management



A statement of English Nature's views about the management of Honeypot Wood, Wendling Site of Special Scientific Interest (SSSI).

This statement represents English Nature's views about the management of the SSSI for nature conservation. This statement sets out, in principle, our views on how the site's special conservation interest can be conserved and enhanced. English Nature has a duty to notify the owners and occupiers of SSSI of its views about the management of the land.

Not all of the management principles will be equally appropriate to all parts of the SSSI. Also, there may be other management activities, additional to our current views, which can be beneficial to the conservation and enhancement of the features of interest.

The management views set out below do not constitute consent for any operation. English Nature's written consent is still required before carrying out any operation likely to damage the features of special interest (see your SSSI notification papers for a list of these operations). English Nature welcomes consultation with owners, occupiers and users of the SSSI to ensure that the management of this site conserves and enhances the features of interest, and to ensure that all necessary prior consents are obtained.

Management Principles

There may be several different ways in which the wood can be managed to best conserve its value for wildlife - by promoting an appropriate woodland structure, by ensuring regeneration and by looking after the things that make this wood special. The attached notes give broad views on a range of regimes that may be appropriate on your site.

A diverse woodland structure with some open space, some areas of dense understorey, and an overstorey of more mature trees (which may be the standard trees under a coppice-with-standards regime) is important. A range of ages and species within and between stands is desirable.

Some dead and decaying wood such as fallen logs, old hollow trees or old coppice stools is essential for providing habitats for fungi and dead wood invertebrates. Work may, however, be needed to make safe dangerous trees where they occur in areas of high public access.

Open spaces, either temporary gaps created by felling or coppicing or more permanent areas such as rides and glades, benefit other groups of invertebrates such as

butterflies. They should be of sufficient size to ensure that sunny conditions prevail for most of the day. Rides and glades may require cutting to keep them open.

Felling, thinning or coppicing may be used to create or maintain variations in the structure of the wood, and non-native trees and shrubs can be removed at this time. To avoid disturbance to breeding birds the work is normally best done between the beginning of August and the end of February. Work should be avoided when the ground is soft, to prevent disturbing the soil and ground flora. Wet woodland by streams and other waterbodies is often best left undisturbed. Normally, successive felling, thinning or coppicing operations should be spread through the wood to avoid too much disturbance in one area. However, where there is open space interest (e.g. rich butterfly populations) adjacent plots may be worked to encourage the spread of species that are only weakly mobile.

Natural regeneration from seed or stump regrowth (as in coppice) is preferred to planting because it helps maintain the local patterns of species and the inherent genetic character of the site.

Deer management and protection from rabbits or livestock are often necessary. Whilst light or intermittent grazing may increase woodland diversity, heavy browsing can damage the ground flora and prevent successful regeneration.

Parts of the wood should be left unmanaged to benefit species that do best under low disturbance. In addition, lack of management allows for the operation of natural processes such as windblow. Within these areas some trees will eventually die naturally and dead wood accumulate.

Where they are a threat to the interest of the wood, invasive introductions such as sycamore should, where practical, be controlled.